

Available online at http://www.journalcra.com

International Journal of Current Research Vol. 5, Issue, 02, pp.017-021, February, 2013

INTERNATIONAL JOURNAL OF CURRENT RESEARCH

# **RESEARCH ARTICLE**

# Variation in seed traits and distribution of *Jatropha curcas* L. in Senegal

### <sup>1\*</sup>Ouattara, B., <sup>2</sup>Diédhiou, I., <sup>2</sup>Ndir, K. N., <sup>1,3</sup> Agbangba, E. C., <sup>1</sup>Cisse, N., <sup>3</sup>Diouf, D., <sup>3</sup>Akpo, E. L. and <sup>4</sup>Zongo, J. D.

<sup>1</sup>Centre d'Etudes Régional pour l'Amélioration de l'Adaptation à la Sécheresse (CERAAS) BP 3320 Route de Khombole Thiès Sénégal

<sup>2</sup>Département de Productions Végétales, Ecole Nationale Supérieure d'Agriculture (ENSA) Université de Thiès BP A296 Route de Khombole Thiès Sénégal

<sup>3</sup>Département de Biologie Végétale, Université Cheikh Anta Diop de Dakar BP 5005 Dakar Sénégal <sup>4</sup>Unité de formation et de recherche en sciences de la vie et de la terre (UFR/SVT), Université de Ouagadougou, 03 BP 7021, Ouagadougou 03, Burkina Faso

ARTICLE INFO	ABSTRACT				
Article History:	A thorough and extensive germplasm exploration survey was undertaken during 2009 and 2010 to assess the				

Received 14<sup>th</sup> November, 2012 Received in revised form 06<sup>th</sup> December, 2012 Accepted 26th January, 2013 Published online 14th February, 2013

Key words:

Jatropha curcas L., Distribution, Seed traits, Senegal

## INTRODUCTION

During the last years, Jatropha curcas L. has gained interest in Sahelian zone as biofuel crop. J. curcas is an oil bearing species and its oil can be converted to biodiesel that meets the American and European biodiesel standards (Tiwari et al., 2007). The interest for J. curcas in Sahelian zone is probably due to its high adaptability to wasteland then can be cultivated on marginal soils without compromising the food, fodder security and improve livelihoods in arid regions (Reddy et al., 2008). It sheds its leaves during the dry season, therefore is well adapted to arid and semi arid conditions characteristic of Sahelian zone. J. curcas belongs to Euphorbiaceae family and native from tropical South America (Carels, 2010). It was introduced in Africa and India by Portuguese seafarers (Heller, 1996). Cultivation of J. curcas can be done on well drained soils and requires low nutrient content (Heller, 1996). However, great variability in J. curcas growth and productivity was reported (Achten, 2008). Annual seed production can range from about 0.2 kg to more than 2 kg per plant (Francis et al., 2005). The major concern is that J. curcas is a wild species that domestication is at embryonic stage. Many authors recommended selection of best genotypes with high yield and oil content and variability studies are carried out to identify genotypes that can be used for amelioration programs of the species. The systematic work on germplasm exploration, characterization, utilization and documentation is of great importance to identify genetic variability for desired traits. Wide variation in 100 seed weight and oil content in accessions collected from India was reported (Ginwal et al., 2005; Rao et al. 2008,

distribution and variability in seed traits of Jatropha curcas L. in Senegal. Nineteen accessions from different agro ecological zones of the country were collected to evaluate variability in seed characters. Trees aged at least 5 years only were considered. Among the seed traits studied, 100 seed weight ranged from 63.68 to 77.83 g and seed length from 17.89 to 19.15 mm. The highest 100 seed weight (77.83 g) was recorded for the accession Jc-16 collected from Mampatim. Accessions from Soudanian zone showed high values of seed traits while low seed traits were recorded in Soudano-Sahelian zone. Variability in seed traits was not linked to geographic location. Old plantations of Jatropha curcas are spread in the central, south and coastal zones of Senegal where the species is well known by the populations. However, the species was least represented above 400 mm isohyet and unfamiliar. No old plantation was observed in extreme north of Senegal.

Copy Right, IJCR, 2013, Academic Journals. All rights reserved.

Kaushik et al., 2007; Wani et al., 2006). In spite of wide spread of J. curcas in Senegal, little work has been done so far in this aspect and information on local J. curcas variability is at infancy stage. Though, tree species with a wide geographical distribution exhibits considerable variation in anatomy, physiology, morphology and genetics to survive and reproduce under varying environmental conditions over generations (Antonovics, 1971; Nienstaedt, 1975). Variation in germination behavior (Ouattara et al., 2011) and seedling growth inoculated with arbuscular mycorhizae (Leye et al., 2009) has been reported in J. curcas accessions from Senegal. Early, Heller (1996) has reported variability in plant growth and seed yield in trial experimentation in Senegal. The study of seed characters of tree born oilseed crops of natural population is often considered to be useful in evaluation of the genetic variability (Mohapatra et al., 2010). Genetic variation in seed traits of J. curcas can be of great potential in tree improvement programs, particularly selection of genotypes having more oil content and yield. The present study aims at identifying the regions of Senegal where old plantations are well represented and characterizing the variability in seed traits of 19 old accessions collected in three agro-ecological zones in the country.

### MATERIAL AND METHODS

Several explorations of germplasm of J. curcas across Senegal have been undertaken between 2009 and 2010. Mature yellow fruits of J. curcas were collected from 19 accessions in November 2010 in three agro-climatic zones of Senegal (Fig 1). Climatic conditions of J. curcas fruits origins were represented in Table 1. Fruits were collected from at least ten trees in the same population spaced at least 10 m and randomly selected to ensure inclusion of local genetic

<sup>\*</sup>Corresponding author: obassiaka@yahoo.fr



Figure 1. Mapping of the 19 accessions of *J. curcas* collected in Groundnut basin, Eastern Senegal and Upper Casamance of Senegal. Seed origins locations in black square; the different colors represent the different agro-ecological zones of Senegal (Map Adapted from Centre national de suivi écologique 1996).

Locality	Seed	Latitude	Longitude	Altitude	Rainfall (mm)	Mean temp. °C
Locality	source code	(°N)	(°W)	(m)	Salack et al. 2011	(min-max)
Maleme thialene	Jc-01	14.04	15.14	31	600-800	35.51-21.73
Hamdalaye pont	Jc-02	13.37	13.35	30	600-800	35.51-21.75
Barkeyel	Jc-03	13.28	13.27	80	600-800	35.51-21.75
Dialacoto	Jc-04	13.18	13.17	61	800-1100	35.51-21.73
Medina Maka	Jc-05	13.13	13.08	62	800-1100	35.51-21.73
Bantancountou	Jc-06	13.15	13.04	52	800-1100	35.51-21.73
Niemenike	Jc-07	12.52	12.21	103	800-1100	34.73-22.2
Mako foukola	Jc-08	12.5	12.21	83	800-1100	34.73-22.2
Bala	Jc-09	14.01	13.09	81	600-800	35.51-21.73
Faskoto	Jc-10	13.54	14.12	58	600-800	35.51-21.73
Djeydina	Jc-11	13.12	15.31	112	800-1100	33.69-20.22
Sare alpha	Jc-12	13.06	15.35	50	800-1100	33.69-20.22
Sefa soukototo	Jc-13	12.49	15.34	52	800-1100	33.69-20.22
Yacine madina	Jc-14	12.46	15.44	30	800-1100	33.69-20.22
Dabo	Jc-15	12.52	14.28	40	800-1100	35.13-20.06
Mampatim	Jc-16	12.53	14.20	64	800-1100	35.13-20.06
Maka	Jc-17	12.50	15.17	21	800-1100	35.13-20.06
Diaroume	Jc-18	12.58	15.37	40	800-1100	35.13-20.06
Ndawene	Jc-19	13.56	14.49	29	600-800	35.51-21.73

variability. In this study, the accession represented fruits collected from trees in the same life fence or from isolated trees in the same farm or village. Trees were at least 5 years old to avoid J. curcas introduced in the country in 2007 from India in order to explore local variability. Accessions were distant each other at least 20 km. Fruits were labeled to keep identity of each accession. The collection sites coordinates were recorded using the Global Positioning System and Arcview 3.3 was used for point mapping. Fruits were air dried until constant weight under similar conditions of temperature and humidity. To determine Seed/fruit ratio, ten samples including 20 fruits each were weighted and seeds from capsule were extracted through manual threshing and weighted. Seed/fruit ratio was determined as the ratio of seeds Weight to the corresponding fruits weight. Three replicates of 100 undamaged seeds of each accession were chosen randomly and their sizes (Length, breadth and thickness) were measured using an electronic vernier calliper. Five random samples including 100 seeds each were taken from each accession to determine seed weight using an electronic scale.

#### Statistical analysis

Variation recorded for each seed trait was partitioned into components due to genetic and environmental factors using Johnson et *al.* (1955) formula. Phenotypic coefficients of variation (PCV) and genotypic coefficients of variation (GCV) were computed as suggested by Burton, (1952). Heritability in Broad sense was calculated according to Allard (1999) as the ratio of the genotypic variance (Vg) to the phenotypic variance (Vp). Genetic advance (GA) as per cent of the mean assuming selection of the superior 5% of the genotypes was estimated in accordance with Johnson et *al.* (1955).

### RESULTS

*J. curcas* mainly occurs in south, central and coastal zones of Senegal. Many old plantations have been observed in the regions of Zinguinchor and Fatick (Fig 2). The species is well known in these localities where it is used as land and farm fences or



Figure 2. Importance of old plantations of *J. curcas* in the different regions of Senegal (the degree of darkness indicated the importance of old plantation of *J. curcas* encountered).

isolated trees in farm or courtyard in villages. *J. curcas* is well known under the name of "*Tabanani*", but in certain villages the plant is called "*Kidi*" or "*Touba taba*". No old plantation above 5 years was found in extreme north and north-east of the country. In certains villages, *J. curcas* oil is used by women in soap production. *J. curcas* was multiplied mainly by cutting to build fences. According to helder met in certain villages, *J. curcas* is a medicinal plant and treats various ills. They highlighted that *J. curcas* leaves are used to treat wounds and toothaches. The latex is used to treat gingivitis affections. Significative differences were recorded among accessions for seed traits (p< 0.01). One hundred Seed weight ranged from 63.68 to 77.83 g with a mean value of 71.42 g. The highest weight (77.83 g) were recorded in Jc-16 from Upper Casamance followed by Jc-10 (76.16 g) from South Groundnut basin (Table 2).

Seed	100 seeds	Length	Breadth	Thickness	Seed/fruit
source	weight (g)	(mm)	(mm)	(mm)	ratio (w/w)
Jc-01	65.03	18.36	10.94	8.84	0.63
Jc-02	70.39	18.55	11.02	8.62	0.71
Jc-03	71.94	18.38	11.13	8.63	0.69
Jc-04	70.9	18.41	11.25	8.64	0.72
Jc-05	73.97	18.82	11.09	8.65	0.73
Jc-06	74.09	18.86	11.02	8.66	0.70
Jc-07	68.89	18.65	11.21	8.61	0.69
Jc-08	72.76	18.63	11.02	8.46	0.71
Jc-09	69.09	18.27	11.15	8.56	0.71
Jc-10	76.16	19.15	11.11	8.61	0.68
Jc-11	72.64	18.86	11.18	8.77	0.71
Jc-12	68.65	18.45	11.02	8.68	0.70
Jc-13	74.97	18.55	11.15	8.76	0.71
Jc-14	69.45	18.59	11.06	8.9	0.69
Jc-15	75.66	18.65	11.01	8.78	0.68
Jc-16	77.83	19.02	11.36	8.91	0.70
Jc-17	69.47	18.69	11.29	8.97	0.72
Jc-18	71.37	18.76	11.24	8.88	0.71
Jc-19	63.68	17.89	10.97	8.45	0.70
SE mean	0.51	0.04	0.02	0.02	0.00

Table 2. Seed traits variability in J. curcas

The lowest 100 seed weight (63.68 g) was recorded in Jc-19 from South Groundnut basin. Mean seed length was 18.61 mm with a maximal of 19.15 mm measured in Jc-10 and minimal of 17.89 mm observed in Jc-19 from Groundnut Basin. Mean seed breadth was 11.12 mm while mean seed thickness was 8.7 mm. In general, high values of seed traits were recorded in Upper Casamance where rainfall closed to 1100 mm. Low seed traits were observed in South Groundnut basin accessions where rainfall was comprised between 600 to 800 mm. However, certain accessions like Jc-10 from South Groundnut basin showed high seed traits. We also noted low seed traits in accession Jc-12 from Upper Casamance. Mean Seed/fruit ratio was 70%. High seed/fruit ratio (0.73) was recorded in Jc-05 and the low value in Jc-01 (0.63). Variability in seed/fruit ratio was weak with approximately similar values in all the accessions. Phenotypic coefficient of variation (PCV) and genotypic coefficient of variation (GCV) were estimated (Table 3). Highest PCV (5.12%) and highest GCV (4.59%) were recorded in seed weight. Low PCV (1.04%) and low GCV (0.96%) were noted in seed breadth. PCV and GCV for seed/fruit ratio were higher than PCV and GCV of all seed traits except in seed weight. The lower heritability in broad sense (76.87%) was recorded in seed/fruit ratio when it was higher than 89.59% for the others seed traits. Genetic advance was higher than 9.45% in seed weight with lowest value (1.97%) observed in seed breadth. Significative correlation has been recorded between seed traits (Table 4). All significative correlation coefficients were positive. Seed length was significatively correlated to all seed traits excepted Seed/fruit ratio. Seed breadth was correlated to all seed traits excepted to 100 Seed weight. Dendrogram with complete linkage based on square euclidean distance between the different traits showed five clusters (Fig 3). Cluster III (green color) and cluster IV (blue color) were morphological closed. Maximum accessions fallen into cluster IV that grouped 9 accessions. Cluster II (yellow color) included only one accession and was closed to cluster I (red color) that grouped 2 accessions.



Figure 3. Dendrogram showing the different clusters based on seed traits of the 19 J. curcas accessions

	Varia	ance	Coefficient of variation (%)		Hanitabilita (based as as)		
Seed source	Phenotypic	Genotypic	Phenotypic	Genotypic	Heritability (broad sense)	Genetic Advance as % mean	
100 seed weight (g)	13.3300	10.7000	5.1254	4.5920	0.8027	9.4596	
Length (mm)	0.0907	0.0830	1.6194	1.5491	0.9151	3.1912	
Breadth (mm)	0.0135	0.0114	1.0470	0.9604	0.8414	1.9784	
Thickness (mm)	0.0218	0.0198	1.6943	1.6162	0.9098	3.3293	
Seed/fruit ratio	0.0004	0.0002	2.7979	2.1507	0.5909	4.4305	
(w/w)							

Table 3. Estimation of genetic variables for	or seed traits in J. curcas
--	-----------------------------

Table 4. Correlation coefficient among seed traits in J. curcas					
Seed trait	Length	Breadth	Thickness	100 Seed weight	
Breadth	0.3654**				
Thickness	0.4214**	0.3961**			
100 Seed weight	0.5930**	0.1917	0.1947		
Seed/fruit ratio	-0.0269	0.2509*	-0.1411	0.1083	
* Significant at the 0.01	level				

\* Significant at the 0.05 level

### DISCUSSION

J. curcas is well known by the populations of south, central and coastal zones of Senegal. Particularly the species is well known in the south Senegal (Casamance) where it seems to be introduced from Bissau Guinea. J. curcas was distributed by Portuguese seafarers from the Caribbean via the Cape Verde Islands and Bissau Guinea to other countries in Africa and Asia (Heller, 1996). According to Berhaut (1967) J. curcas is known under various names as "Tabanani", "Kidi" in Senegal. However, the author didn't mentioned the name "Touba taba". Based on old plantations distribution in Senegal, Casamance, Eastern Senegal and South Groundnut basin zones are the favorable areas where J. curcas establishment would be easy. Since J. curcas is well known and occurring in these zones as live fences in land or farm, it is adapted to their environmental conditions and populations will adopt easily the High variability was observed in J. curcas seed traits species. among the accessions. 100 seed weight ranged from 63.68 to 77.83 g that is in line with Halilu et al. (2011) who reported that 100 seed weight in their samples ranged from 28.558 to 80.046 g. Variability in environmental conditions of mother trees may explain the variation observed in seed traits. Indeed, J. curcas accessions were collected in different agro ecological zones that present different climatic conditions and soil characteristics. Soil and climate of the place of seed origin are considered as important factors affecting the seed traits (Salazar and Quesada, 1987). Such variations in habitat have been reported in J. curcas (Kaushik et al., 2007, Ginwal et al., 2005) and others tree species (Jindal et al., 1999; Gera et al. 1976, Kumar et al., 2004).

According to Mathur et al. (1984), variability occurring in seed traits might be attributed to genetic in nature as a result of adaptation to various environmental conditions prevailing throughout their distributional range. However, environmental conditions cannot explain alone all the variability observed in seed traits. Accessions from the same agro ecological zone often presented various seed traits. It is clear that one part of the variability is under genetic control. Some genotypes of J. curcas seem to produce large seed traits than others. This is confirmed by the few difference between phenotypic variance and genotypic variance. High values of heritability also indicated that variability in seed traits had genetic base. All the seed traits except seed/fruit ratio showed heritability superior to 80%. The important place of heritability in plant improvement program as it provides an index of the relative role of heredity and environment in the expression of various traits has been reported (Kaushik et al., 2007). Comparing seed traits variability it has been noted that 100 seed weight recorded the highest variability. Genetic advance as percent of mean was high for 100 seed weight (9.45%) further suggested the potentiality of the test material for future improvement through selection. High heritability and high genetic advance for 100 seed weight have been reported by Kaushik et al. (2007) in J. curcas. Hence these characters can be considered

as best gain characteristics for *J. curcas* improvement program because of its strong genetic control and the wide variability. In addition, in early reports (Ponnammal *et al.*, 1993, Kaushik *et al.*, 2001; Kaushik *et al.*, 2003), high seed traits recorded better germination and seedling growing. Halilu *et al.* (2011) highlighted that 100 seed weight is positively correlated to oil content. So, attention can be done to accessions with high seed traits that could be useful for breeding programs. Positive correlation coefficients were recorded between seed traits. Seed weight was correlated to seed length. Similar results were found by Mohapatra and Panda (2010) who reported positive correlation between seed length and 100 seed weight. Contrary to Kaushik *et al.* (2007), no significance correlation has been found between 100 seed thickness.

Seed length was found positively correlated to seed breadth and seed thickness. Similar results have been reported in Acacia catechu (Ramchandra et al., 1996; Kumar et al., 2004). Strong positive correlation among seed traits indicated that genes involved in these traits might be closely linked or might have pleitropic effects (Das et al., 2010). The dendrogram based on seed traits fallen to group accessions from the same agro ecological zone together. Indeed, accessions from Upper Casamance and South of Eastern Senegal are grouped together. So geographical diversity did not necessarily represent genetic diversity. Similar results have been reported in J. curcas (Kaushik et al., 2007) and C. tetragonaloba (Saini et al., 2004). However, at the level of 45 of similarity, two majors cluster were formed. The first cluster grouped accessions from Soudanosahelian zone, were rainfall varies between 500 to 900 mm, and the second cluster grouped accessions from Soudanian zone, where rainfall varies from 900 to 1100 mm (FAO, 2007). Cluster from Soudanian zone showed large seed traits. Wet zones are most favorable to J. curcas and the species is not common in regions with arid and semi-arid climate (Maes et al., 2009). As described by several authors (Kaushik et al., 2007; Ginwal et al., 2005), variation in J. curcas seed sources with respect to their morphological characters could be due to fact that the species grows over a wide range of rainfall, temperature and soil type.

#### Conclusion

The present study revealed that considerable genetic variability exists in *J. curcas* in Senegal with respect to seed morphology. On the basis of our results, wet zones are favorable to *J. curcas* where it produces high values of seed traits. *J. curcas* is widely distributed in south and central of Senegal where the species is well known. Accessions with high 100 seed traits identify hereby might be used in further breeding programs. This study is merely an indication that variability in seed traits is present in senegal and researches are need to identify consistently high yielding and high oil content in local ecotypes that are best adapted to Senegal climatic conditions.

#### Acknowledgements

The authors thank RIPIECSA for financial support. OUATTARA thanks DAAD for scholarship award. The authors also are thankful to Prof. Naresh KUMAR for statistic documentations.

#### REFERENCES

- Achten W.M.J., Verchot L., Franken Y.J., Mathijs E., Singh V.P., Aerts R., Muys B., 2008: J. curcas bio-diesel production and use. Biomass and bioenergy 32 1063-1084.
- Antonovics J., 1971: The effects of a heterogenous environment on the genetics of natural populations. Am Sci 59(5): 593–595.
- Berhaut J., 1967 Flore du Sénégal. 2éme édition. Dakar. Librairie clairafrique. 485p.
- Burton GW, 1952: Quantitative inheritance in grass. Proceedings of the sixth. Silvae Genetica 1952;37:221–7.
- Carels N., 2010: J. curcas curcas: A Review. Advances in Botanical Research, Vol. 50. DOI: 10.1016/S0065-2296(08)00802-1
- Das S., Misra R.C., Mahapatra A.K., Gantayat B.P. and Pattnaik R.K. 2010: Genetic Variability, Character Association and Path Analysis in *J. curcas curcas*. World Applied Sciences Journal 8 (11): 1304-1308.
- FAO, 2007: Floods caused damage in most countries but increased rains in august improve crop prospects. Sahel Weather and Crop Situation Report N° 3.
- Francis, G., Edinger, R. and Becker, K. A., 2005: A concept for simultaneous wasteland reclamation, fuel production, and socioeconomic development in degraded areas in India: Need, potential and perspectives of J. curcas plantations. Natural Resources Forum 29, 12–24.
- Gera N, Gera M, Purohit M, 1976: Source variation in seed and germination characteristics in Acacia nilotica ex Del. Seed Research 28:27–31.
- Ginwal, H. S., S. S. Phartyal, P. S. Rawat and R. L. Srivastava 2005: Seed source variation in morphology, germination and seedling growth of J. curcas curcas Linn. in Central India. Silvae Genetica 54:76-80.
- Heller Joachim, 1996. Physic nut. J. curcas curcas L. Promoting the conservation and use of underutilized and neglected crops. Institute of Plant Genetics and Crop Plant Research Notes, Gatersleben/ International Plant Genetic Resources Institute, Rome Italy, pp 66.
- Jindal SK, Satyavir, Pancholy A, 1999: Variability and associations for seed yield, oil content and tree morphological traits in neem (Azadirachta indica). Journal of Tropical Forest Science 1999;11:320–2 [special 10th year anniversary issue].
- Johnson, H. W., H. F. Robinson and R. F. Comstock, 1955: Estimates of genetic and environmental variability in Soyabean. Agron. J. 47: 314–318.
- Kaushik N., 2001: Effect of seed traits on the performance of top feed tree species at seedling stage. Forage Research 27:43–5.
- Kaushik, N., J. C. Kaushik and S. Kumar, 2003: Response of J. curcas curcas to seed traits and growing medium. Journal of Non-Timber Forest Products 10:40-2.
- Kaushik, N., K. Kumar, S. Kumar and S. Roy, 2007: Genetic variability and divergence studies in seed traits and oil content of J. curcas (J. curcas curcas L.) accessions. Biomass and Bioenergy 31:497-502.

- Kumar R, Nautiyal S, Kumar P, Bahuguna A, 2004: Seed source variation in Khair (Acacia catechu Willd.). Indian Forester 130:530–6.
- Leye E.H.M., Ndiaye M., Ndiaye F., Diallo B., Sarr A.S., Diouf M. et Diop T., 2009 : Effet de la mycorhization sur la croissance et le développement de *J. curcas curcas* L. *Revue des Energies Renouvelables Vol. 12 N°2 (2009) 269 – 278*
- Maes W.H., Trabuccoa A., Achten W.M.J., Muysa B.; 2009: Climatic growing conditions of J. curcas curcas L. biomass and bioenergy 33 1481-1485.
- Mathur, R. S., K. K. Sharma and M. M. S. Rawat, 1984: Germination behaviour of provenances of *Acacia nilotica* sp. indica. Indian Forester 110: 435–449.
- Nienstaedt H., 1975: Adaptive variation Manifestations in tree species and uses in forest management and tree improvement. Proc. 15th Can. Tree Improv. Assoc. Part 2, pp. 11–23.
- Ouattara B., Diédhiou I., Ndoye K. Diouf D., Akpo E. L., 2011: Effect of water regimes and pre-sowing treatments on seeds germination of different provenances of J. curcas curcas L. in Senegal. International Journal of Science and Advanced Technology. Vol: 1(9), 151-156.
- Ponnammal NR, Arjunan MC, Antony KA, 1993. Seedling growth and biomass production in Hardwickia binnata Roxb as effected by seed traits. Indian Forester 119:59–62.
- Ramchandra NG, 1996: Provenance variation in seed and seedling parameters in Acacia catechu Willd. PhD thesis, FRI Deemed University, Dehra Dun, 1996.
- Reddy B V S, Ramesh S, Ashok Kumar A, Wani S P, Ortiz R and Ceballos H. 2008: Bio-fuel crops research for energy security and rural development in developing countries. *Bioenergy Resource* 1: 248-258.
- Sagar Mohapatra and Prasanna Kumar Panda, 2010: Genetic Variability on Growth, Phenological and Seed Characteristics of J. curcas curcas L. Notulae Scientia Biologicae 2 (2) 2010, 127-132
- Saini ML, Jain P, Singh JV, 2004. Genetic diversity in a germplasm collection of guar [Cyamopsis tetragonaloba (L.) Taub.]. Forage Research 30:92–5.
- Salack S., Muller B., and Gaye A. T.; 2011: Rain-based factors of high agricultural impacts over Senegal. Part I: integration of local to sub-regional trends and variability. Theor Appl Climatol DOI 10.1007/s00704-011-0414-z
- Salazar, R. and M. Quesada, 1987: Provenance variation in *Guazuma ulmifolia* L. in Costa rica. Commonwealth Forestry Review 66, 317–324.
- Tefera, H., K. Assefa, F. Hundera, T. Kefyalew and T. Teferra, 2003: Heritability and genetic advance in recombinant inbred lines of tef (*Eragrostis tef*). Euphytica 131:91-96.
- Tiwari, A. K., A. Kumar and H. Raheman, 2007: Biodiesel production from J. curcas (*J. curcas curcas*) with high free fatty acids: an optimized process. Biomass and Bioenergy 31:569-75.
- Wani S P, Osman M, D'silva E and Sreedevi T K. 2006: Improved livelihoods and environmental protection through biodiesel plantations in Asia. Asian Biotechnology Develop Review 8(2): 11-29.

\*\*\*\*\*\*